


-23-

CLAIMS


1. A process for the preparation of ZD6126 Phenol:



5

ZD6126 Phenol

from a ZD6126 Alcohol of formula (II):




(II)

wherein R^2 are each independently hydrogen, C_{1-4} alkyl or aryl which comprises:


- 10 reacting said ZD6126 Alcohol of formula (II) with an acid catalyst and an oxidising agent.

2. A process according to claim 1 wherein the acid catalyst is a sulfonic acid.
3. A process according to claim 1 wherein the acid catalyst is methanesulfonic acid.
- 15 4. A process according to any one of the preceding claims wherein the reaction is carried out in the presence of a solvent selected from an aromatic solvent, an ester and an ether.
5. A process according to any one of claims 1 to 3 wherein the reaction is carried out in
- 20 an aromatic solvent selected from toluene and chlorobenzene, or a mixture of two or more of said solvents.
6. A process for the preparation of ZD6126 Phenol:

-24-


**ZD6126 Phenol**

from an allocolchicine or an ester derivative thereof of formula (I):

**(I)**

wherein R^1 is hydrogen, C_{1-6} alkyl or aryl; which comprises:

a) reacting said allocolchicine or an ester derivative thereof of formula (I) with a suitable organometallic reagent and / or a suitable reducing agent; in one or more ethereal solvents to form ZD6126 Alcohol of formula (II):

**(II)**

wherein R^2 is hydrogen, C_{1-4} alkyl or aryl; and

b) reacting ZD6126 Alcohol of formula (II) with an acid catalyst and an oxidising agent.

7. A process according to claim 6 wherein R^1 is C_{1-4} alkyl or aryl.

8. A process according to claim 6 wherein in step a) of the process the allocolchicine or an ester derivative thereof of formula (I) is reacted with a suitable organometallic reagent and wherein R^1 is C_{1-4} alkyl or aryl.

5

9. A process according to any one of claims 6 to 8 wherein the organometallic reagent in step a) of the process is selected from a compound of the formula R^2-X , wherein R^2 is as defined claim 6 and X is a magnesium halide or lithium.

10 10. A process according to any one of claims 6 to 8 wherein the organometallic reagent in step a) is methyllithium.

11. A process according to any one of claims 6 to 10 wherein the one or more ethereal solvents is selected from tetrahydrofuran, diethyl ether, diethoxymethane, 2-ethoxyethylether,
15 2-methoxyethyl ether and dimethoxy ethane, or a mixture of one or more of said solvents.

12. A process any one of claims 6 to 11 wherein in step a) the allocolchicine or an ester derivative thereof of formula (I) is added to a reaction mixture comprising the organometallic reagent.

20

13. A process according to claim 12 wherein the organometallic reagent is methyllithium.

14. A process according to any one of claims 6 to 13 wherein the acid catalyst in step b) is a sulfonic acid.


25

15. A process according to claim 14 wherein the acid catalyst in step b) is methanesulfonic acid.

16. A process according to any one of claims 6 to 15 wherein in step b) of the process is
30 carried out in the presence of a solvent selected from an aromatic solvent, an ester and an ether.

-26-

17. A process according to any one of claims 6 to 15 wherein in step b) of the process is carried out in the presence of an aromatic solvent selected from toluene and chlorobenzene, or a mixture of two or more of said solvents.
- 5 18. A process according to any one of claims 6 to 17 wherein, the process is effected in one stage, without isolation of ZD6126 Alcohol of formula (II).
19. A process according to any one of claims 6 to 18 wherein R^1 is C_{1-4} alkyl.
- 10 20. A ZD6126 Alcohol of formula (II) as defined in Claim 1, with the proviso that R^2 cannot both be methyl or both be hydrogen.
21. A process for the preparation of a ZD6126 Alcohol of the formula (II) as defined in claim 6 which comprises reacting allocolchicine or an ester derivative thereof of the formula (I)
- 15 as defined in claim 6 with a suitable organometallic reagent and/or suitable reducing agent in one or more ethereal solvents.
22. Use of a ZD6126 Alcohol of formula (II) as defined in claim 1 in a process for the preparation of ZD6126 Phenol.
- 20
23. A ZD6126 Alkene of formula (III):



(III)


wherein R^2 is hydrogen, C_{1-4} alkyl or aryl and R^3 is hydrogen or C_{1-3} alkyl.

-27-

24. A process for the preparation of ZD6126 Alkene of formula (III) as defined in claim 23 which comprises reacting a ZD6126 Alcohol of the formula (II) as defined in claim 1 wherein at least one R^2 group is C_{1-4} alkyl, with an acid catalyst.

5 25. A process for the preparation of a ZD6126 Phenol which comprises reacting a ZD6126 Alkene of formula (III) as defined in claim 23 with an acid catalyst and an oxidising agent.

26. A ZD6126 Hydroperoxide of formula (IV):




(IV)

wherein R^2 are each independently hydrogen, C_{1-4} alkyl or aryl.

27. A process for the preparation of a ZD6126 Hydroperoxide of formula (IV) as defined in claim 26 which comprises reacting a ZD6126 Alcohol of the formula (II) as defined in claim 1 with an acid catalyst and oxidising agent.

28. A process for the preparation of ZD6126 Phenol which comprises reacting a ZD6126 Hydroperoxide of formula (IV) as defined in claim 26 with an acid catalyst.

29. A ZD6126 Reactive Dimer of formula (V):



-28-

(V)

wherein R^2 are each independently hydrogen, C_{1-4} alkyl or aryl.